

Research article

Differences between wild type and mutant isolate of *Serratia* by the amount of prodigiosin extracted and antibacterial activity against different types of bacteria

Hussam Mahmood Hasan Al-imari^{1*}

ABSTRACT

The mutation was applied on *Serratia marcescens* (*S. marcescens*) to make it produce prodigiosin with higher amount. One isolate of *S. marcescens* (*Wild*) was exposed to different kinds of antibiotics (Gentamycin 10 µg/ml) to produce mutant isolates from *S. marcescens* and examined to produce prodigiosin. The isolate that produced the largest amount of prodigiosin among four isolates was used for further studies. The antibacterial activity of prodigiosin that produced from bacterial strains was examined against different species of bacteria, *Staphylococcus*, *Bacillus*, *E. coli*, and *Pseudomonas*.

Keywords: Antibacterial activity, Prodigiosin, *Serratia marcescens*.

Citation: Al-imari HMH. (2016) Differences between wild type and mutant isolate of *Serratia* by the amount of prodigiosin extracted and antibacterial activity against different types of bacteria. *World J Exp Biosci* 4:15- 19.

Received November 24, 2015; Accepted December 28, 2015; Published January 8, 2016.

INTRODUCTION

Prodigiosin (red pigment) is a member of prodigiosins produced by *Serratia*, actinomycetes and few other bacteria. Different concentrations of prodigiosin were studied against cancerous cells. It was involved in apoptosis of haematopoietic cancer cell [2]. *Serratia* is a Gram negative bacteria, opportunistic pathogen for human, plant and insect and member of enterobacteriaceae. Strains of *S. marcescens* produce nosocomial

mial infections that are clinically problematic because multidrug resistance. *S. marcescens* has been isolated from different niches (soil, water, plants and air), *S. marcescens* is a non-motile, citrate positive bacterium. On nutrient agar, the colonies are convex, circular with entire margin [2]. Some *S. marcescens* strains were able to grow in a wide range of ecological niches has been attached to the protection of a spectrum of extracellular



*Correspondence: Al-imari HMH.hussamalammar@yahoo.com.
Department of Biology, College of Science, University of Baghdad, Baghdad, Iraq.
Full list of author information is available at the end of the article.

Copyright: © 2016, Al-imari HMH. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any site, provided the original author and source are credited.

Al-imari HMH. (2016).

products including bacteriocins, chitinase, proteases, nucleases, lipases, and surfactants, wetting agents [3]. A bifurcated pathway has been proposed for the productions of prodigiosin culminating in the enzymic condensation of the final products of the two pathways, 4-methoxy-2,2-bipyrrole-5-carboxyaldehyde (MBC) and the monopyrrole, 2-methyl-3-n-amylyl-pyrrole (MAP). The precursors for prodigiosin were seemed to be acetate, serine, alanine, methionine and proline [4]. Recently, the mechanism of proline incorporation into a pyrrole moiety has been shown biochemically and a pathway for production of undecylprodigiosin proposed [3]. Prodigiosin have no special role in the physiology of strains, but have been reported to have antibacterial, antifungal and antiprotozoal/antimalarial activities [5].

They are of interest because they might have critical clinical utility. Prodigiosins and synthetic derivatives have been shown to have main and specific immunosuppressive activity, with novel targets clearly distinct from other drugs [6]. Many possible mechanisms are suggested attributed to prodigiosins as pH modulators, cell cycle inhibitors, DNA cleavage agents and mitogen activated protein kinase regulators [7]. These molecules when combined with some other anticancer agents can help in fighting cancer [8]. The immunosuppressive characters were also carried earlier by Han et al. (1998) [9].

Effect of prodigiosin on human carcinoma cells was tested by Kamble, et al. (2012), and significant results were found. *Spodoptera litura* is classified as one of the crop destructing insect. *Bacillus thuringiensis* toxin viz Cry1C is the major weapon used against these insects, insecticidal activity was found to be increased when prodigiosin was combined with Cry1C [8]. For the reasons above, the importance of prodigiosin was revealed. Therefore, the amount of prodigiosin was hoped to be increased by doing spontaneous mutation. In present study the antibacterial effect of prodigiosin was evaluated.

MATERIALS AND METHODS

Selected isolates

Environmental isolates were collected, examined for growing of *Serratia*, many isolates were identified as *S. marcescens* that had the ability to produce the red pigment (prodigiosin) (Fig 1).

Extraction of prodigiosin

The isolates were examined for production of prodigiosin to select the best isolate in prodigiosin production. The best isolate that produce prodigiosin in large amount was called **wild**.

Nutrient agar antibiotic medium for gradient method [10,12]

The nutrient agar was prepared as per the manufacturer's instructions (Himedia, India). The

medium was mixed with antibiotic solution at 40 °C and poured in sterilized plates (for gradient plates and selective plates) and incubated for 24 h to insure the sterility [10-12].




Fig 1. *S. marcescens* isolate wild produced prodigiosin attributed with red pigment.

This medium was used for isolation bacterial mutant of different antibiotics. Two hundred of overnight broth medium was transferred to nutrient agar gradient plates, spread by sterilized spreader carefully, Incubated at 37°C for 24 h. The growth of bacteria was examined according to action of antibiotic concentration on gradient plates (Fig 2).



Fig 2. Gradient plate inoculated with *S. marcescens* and incubated for 37°C for 24 h.

Production of spontaneous mutant isolates

The selected isolate of *S. marcescens*, **wild** isolate was exposed to various concentrations of different types of antibiotics to produce mutated isolates. Two hundred of nutrient broth culture (overnight 18 h) was transferred to nutrient agar antibiotic plates for different antibiotic with different concentrations (according the estimation of

Al-imari HMH. (2016).

active concentration by gradient method, spread by sterilized spreader, then incubated at 37°C for 24 h. The number of mutants appeared colonies were counted. Small part of selected colonies were streaked by sterilized loop on the same nutrient agar antibiotic medium, which the colonies were selected from it to be sure these colonies were mutant and not physiological adaptation compared with type strain, the colony which failed to grow was to be not real mutant.

Extraction of prodigiosin pigment:

Three of mutant isolates G1-4, G1-2, G0.5-4, and the wild isolate were tested and the comparison among them was done according to the amount of prodigiosin pigment produced.

Antimicrobial activity of prodigiosin:

Four isolates, *Pseudomonas*, *Staphylococcus*, *Bacillus*, *Escherisea coli* were tested for susceptibility against prodigiosin pigment.

Statistical analysis

Statistical software SPSS 10.0 was applied in this experiment. Significance between produced prodigiosin by selected isolate and other mutant isolates was estimated.

RESULTS

Two isolates were identified as *S. marcescens*, one of the two isolates was been able to produce prodigiosin in perceptible amount called **wild**, while another one was not been able to produce prodigiosin in perceptible amount, the second isolate was neglected. After purification of **wild** isolate, it was examined again for its ability to produce prodigiosin. After that it was exposed to solution of different kinds of antibiotics (gentamycin, ceftriaxone, cefalexin) with different concentrations, gentamycin was the only antibiotic was not decrease or prevent production of prodigiosin. In **fig 3** the growth that exposed to cefataxime showed no prodigiosin production, while the plate that exposed to gentamycin was not effected on the prodigiosin production.




Fig 3. Isolate of *S. marcescens* exposed to ceftriaxone (a) and gentamycin (b).

Gentamycin antibiotic was mixed with nutrient agar media to produce spontaneous mutants from wild (**Fig 4**).




Fig 4. Different mutant isolates of *Serratia* grown on medium contains gentamycin.

Many mutant isolates were isolated from wild isolate (**Fig. 4**), and examined for prodigiosin production. Three mutant isolates shown in table 1. Production of prodigiosin in in mutant isolates was higher than wild isolate.

Table 1. Production of prodigiosin from *wild* and mutant isolates.

isolates	Gentamycin concentration (mg/ml)	Amount of produced prodigiosin (µg)
wild	Without	150000
G1-2	1	280000
G1-4	1	190000
G0.5-4	0.5	250000

There is significant difference ($P < 0.05$) in production of prodigiosin by wild isolate and three mutant isolates, while there is no significant differences ($P < 0.05$) in comparison with other mutant isolates. An application of produced prodigiosin by wild isolate and other three mutant isolates was examined. This application was antibacterial activity of prodigiosin produced by wild isolate and the three mutant isolates against different Gram positive and negative bacteria (**Table 2**).

Table 2. Antibacterial activity of prodigiosin of *wild* isolate and three mutant isolates against *Pseudomonas*, *Staphylococcus*, *Bacillus* and *E. coli*.

	<i>Pseudomonas</i>	<i>Staphylococcus</i>	<i>Bacillus</i>	<i>E. coli</i>
Wild	9mm	10mm	9mm	7mm
G1-2	9mm	11mm	10mm	7mm
G1-4	9mm	10mm	8mm	8mm
G0.5-4	8mm	8mm	16mm	8mm
Control	-	-	-	-

The results in **table 2** showed that the antibacterial activity of prodigiosin produced from wild isolate and three mutant isolates against different Gram positive and

negative bacteria, and revealed that prodigiosin pigment (whether extracted from wild isolate or from mutant isolates) had antibacterial activity against both Gram positive and negative bacteria. **Fig. 5** showed significant effect ($P < 0.05$) at the concentration of prodigiosin (1000 $\mu\text{g/ml}$).




Fig 5. Effect of prodigiosin pigment of wild and other mutants isolates of *S. marcescens* on growth of different bacteria, *Pseudomonas*, *Staphylococcus*, *Bacillus* and *E. coli*.

Prodigiosin pigment had inhibition activity against different bacteria isolates with concentration of 1000 $\mu\text{g/ml}$.

DISCUSSION

In present study, two isolates were purified and identified as *S. marcescens*, **wild** isolate was chosen because it produces prodigiosin in significant amount. While another isolate was neglected because it could not be able to produce prodigiosin and this disagrees with the aim of this study, (production prodigiosin pigment in large amounts). The result of gradient plates was showed that **wild** isolate grew on low concentrations of antibiotic (gentamycin, ceftriaxone and cefalexin), gentamycin was showed best results in comparing with other two antibiotics as shown in **figs 3** and **4**. The **wild** isolate succeeded to grow on the regions of low concentrations of antibiotics, while failed to grow on the regions of high concentrations. That means the bacteria, **wild** isolate could resist the low concentrations of antibiotics and survival again and again because mutations was happened for this bacteria make it resist the presence of antibiotics that must kill it, these isolates was called **spontaneous mutants**. While, the death of isolates on media with antibiotics when still inoculated on the same media with antibiotics that's mean there is no real mutation and the bacteria adapt the low concentrations in the first time but could not still growing in the presence of this antibiotic. The results were showed that the amount of prodigiosin produced by **wild**

isolate increased after spontaneous mutation (**Table 1**). The activity of prodigiosin that produced in largest amount by mutant isolate had not changed their activity after applying spontaneous mutation, the production of prodigiosin from *S. marcescens* that produced in liquid media was more than amount produced by solid media, and this was agreed with previous study [13]. The reason of production large amount of prodigiosin after obtaining mutant isolates, is the spontaneous mutant isolate was stronger than wild isolate and had more able to produce prodigiosin [12].

The red pigment was showed a serious role in inhibition the growth of bacteria, and this was agreed with earlier studies [13,7]. The method of inhibition of prodigiosin was returned to considering prodigiosin as cell cycle inhibitors, and DNA cleavage agents [1]. The amount of prodigiosin was increased because the great role of prodigiosin as one of new experimented way to fight cancer, effect of prodigiosin on human carcinoma cells was investigated [14, 9].

Prodigiosin combined with another materials to enhance its activity, like combination between toxin viz Cry1C that was extracted from *B. thuringiensis* and prodigiosin, this combination is the major weapon employed against *Spodoptera litura*, that is one of the crop destructing insect, the insecticidal activity was found to be enhanced when prodigiosin was combined with Cry1C. [14-16]. The present study proved strongly the direct role of prodigiosin in inhibiting bacterial growth.

Conflict of interest

The authors declare that they have no conflict of interests.

REFERENCE

- Richardson WS, Wilson MC, Nishikawa J, Hayward RS. (1995). The well-built clinical question: a key to evidence-based decisions. *ACP J Club* **123**: A12-13.
- American Dental Association. (2008) Policy on evidence-based dentistry: introduction. <http://www.ada.org/1754.aspx>. Accessed 8 Mar 2015.
- Gray GE, Pinson LA. (2003) Evidence-based medicine and psychiatric practice. *Psychiatr Q* **74**:387-99.
- Prisant LM. (1997) Hypertension In: Conn RB, Borer WZ, Snyder JW, editors. *Current Diagnosis*. Philadelphia: W.B. Saunders: 349-59.
- Sackett DL, Rosenberg WM, Gray JA, Haynes RB, Richardson WS. (1996) Evidence based medicine: what it is and what it isn't. *Brit Med J* **312**:71-72.
- Brownson RC, Baker EA, Leet TL, Gillespie KN, True WN. (2003) *Evidence-Based Public Health*. New York: Oxford University Press.
- McQuay HJ, Moore RA. (1988) *Evidence-based resource for pain relief*. Oxford:Oxford University Press.
- Rosenberg W, Donald A. (1995) Evidence based medicine: an approach to clinical problem solving. *Brit Med J* **310**:1122-26.
- Burls A. (2009) What Is Critical Appraisal? London: Hayward Group. http://www.medicine.ox.ac.uk/bandolier/painres/download/whatis/what_is_critical_appraisal.pdf. Accessed 14 Mar 2015.
- Attia A. (2005) Bias in RCTs: confounders, selection bias and allocation concealment. *Middle East Fertil Soc J* **3**:258-61.
- Sackett DL. (1997) Evidence-based medicine. *Semin Perinatol* **21**:3-5.
- Atkins D, Best D, Briss PA, Eccles M, Falck-Ytter Y, et al. (2004) GRADE Working Group. Grading quality of evidence and strength of recommendations. *Brit Med J* **328**:1490.
- Schunemann HJ, Hill SR, Kakad M, Vist GE, Bellamy R, et al.

- (2007) Transparent development of the WHO rapid advice guidelines. *PLoS Med* 4:e119.
14. **Brozek JL, Akl EA, Alonso-Coello P, Lang D, Jaeschke R, et al.** (2009) GRADE Working Group. Grading quality of evidence and strength of recommendations in clinical practice guidelines. Part 1 of 3. An overview of the GRADE approach and grading quality of evidence about interventions. *Allergy* 64:669–77.
15. **Kiriakou J, Pandis N, Madianos P, Polychronopoulou A.** (2014) Developing evidence-based dentistry skills: how to interpret randomized clinical trials and systematic reviews. *Prog Orthod* 51-58.
-

Author affiliation:

1. *Department of Biology, College of Science, University of Baghdad, Baghdad, Iraq.*

